

Multiple Scattering And Volume-Roughness Interactions In Sea Bed Acoustics

Anatoliy N. Ivakin
Andreev Acoustics Institute
Shvernika 4, Moscow 117036, Russia
email: aniva@glasnet.ru

Darrell R. Jackson
Applied Physics Laboratory,
College of Ocean and Fishery Sciences,
University of Washington, Seattle, Washington 98105
phone: (206) 543-1359, fax: (206) 543-6785, email: drj@apl.washington.edu
Award Number: N00014-97-1-0615

LONG-TERM GOAL

The objective of this research is to improve understanding of bottom acoustic scattering at mid- and high-frequencies and to improve conventional algorithms for remote acoustic characterization of marine sediments.

OBJECTIVES

The primary scientific objective is to develop improved theoretical models for seabed scattering using approximations having a wider range of applicability than those presently available.

APPROACH

Scattering mechanisms due to volume inhomogeneities of the sediment and roughness of its surface and internal interfaces, as well as volume-roughness interactions are considered in the frame of a unified approach [1]. Average seabed parameters can be arbitrary functions of depth and are included in the zeroth-order solution for the field and Green's function. A first-order iteration of the unified integral equation gives the first-order perturbation solution in a form more convenient for describing scattering from sea beds with complex structure. A generalization for the case of stratified fluid sediments covering rough inhomogeneous elastic half-space (basement) can be done as well [1-3].

The unified integral equation method can permit in principle a rigorous consideration of scattering in the complex seabed medium with strong volume and roughness irregularities. However, because a direct numerical approach entails a large computational burden for realistic medium parameters, the main practical application of the unified approach in the near future is validation of various analytical approximate methods considering multiple scattering effects. A Cumulative Forward-scatter Single-Backscattering (CFSB) approximation [de Wolf, 1971] seems to be promising and its modification can be applicable for treating multiple scattering in sea bed sediments [4].

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1999		2. REPORT TYPE		3. DATES COVERED 00-00-1999 to 00-00-1999	
4. TITLE AND SUBTITLE Multiple Scattering And Volume-Roughness Interactions In Sea Bed Acoustics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 NE 40th Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

WORK COMPLETED

The general first-order model developed previously has been used to analyze the frequency-angular dependencies of the scattering strength for both monostatic and bistatic cases for various seabed types at mid- and high-frequencies. Work has continued on the problem of acoustic classification of sediments, including the sensitivity to surficial properties and the signatures of sediment roughness and volume in spatial correlation of the backscattered field.

RESULTS

The first-order analysis shows a number of distinctive features in scattering that in turn can be used as classification clues for determination of various parameters of the seafloor. Additional clues are revealed from an analysis of the spatial correlation function of the scattered field that can be used for distinguishing and/or separating the volume and roughness components of scattering [5,6].

Continued model-data comparisons using data from the CBBL experiment at Key West have shown the importance of gradients of the acoustical parameters in the thin (6 cm) top layer of sediments on roughness scattering for both monostatic and bistatic configurations [7]. Also, the sensitivity of measurable scattering characteristics (e.g., the scattering strength and its angular-frequency dependencies) is shown to be extremely strong with respect to surficial parameters. Therefore, determining the parameters at the sea bed surface by various methods (including non acoustical and in situ) is critical for providing the environmental data base for underwater acoustics experiments and those parameters can be considered as classification clues for remote acoustic characterization of marine sediments.

IMPACT/APPLICATION

The models of seabed scattering developed in this research will provide a better understanding of bottom acoustic interaction at mid- and high-frequencies and can be used as a basis for improved algorithms for remote acoustic inversions for seafloor properties.

TRANSITIONS

The results of this work are being adapted in practical models for seabed scattering. For example, a high-frequency bistatic scattering model funded by the ONR Torpedo Environments Program (6.2) incorporates the elastic scattering model developed as part of this work. The correlation method for identifying and/or separating the volume and roughness components of scattering is proposed for use in ASIAEX and other ONR experiments.

RELATED PROJECTS

This research is conducted jointly with the separately funded work of D.R. Jackson and comparisons with CBBL data have been carried out in collaboration with Kevin Briggs, Kevin Williams, Chris Jones and Tim Orsi. The approaches and models developed in this research are relevant to acoustic penetration and multiple scattering issues arising within the ONR Departmental Research Initiative on high-frequency sound interaction with the seafloor.

REFERENCES

1. Ivakin, A.N., 1998. "A unified approach to volume and roughness scattering," J. Acoust. Soc. Am., 103, 827-837.
2. Jackson D.R. and A.N. Ivakin, 1998, "Scattering from elastic sea beds: First-order theory," J. Acoust. Soc. Am., 103, 336-345.
3. Ivakin, A.N. and D.R. Jackson, 1998, "Effects of shear elasticity on sea bedscattering: Numerical examples," J. Acoust. Soc. Am., 103, 346-354.
4. A.N. Ivakin, 1999, "Multiple scattering in marine sediments with volume inhomogeneities", J. Acoust. Soc. Am., 104, 138th Meeting of Acoust. Soc. Amer., Columbus, OH.
5. A.N. Ivakin, 1999, "Seafloor scattering: Modeling and analysis of classification clues", J. Acoust. Soc. Am., 103, 137th Meeting of Acoust. Soc. Amer., Berlin.
6. A.N. Ivakin, 1999, "Seabed volume and roughness scattering: Models and data analysis", ONR Shallow Water Acoustics workshop, Santa Fe, NM.
7. K.Briggs, K.Williams, D.Jackson, C.Jones, A.Ivakin and T.Orsi, 1999, "Influence of fine-scale sedimentary structure on high-frequency acoustic scattering", Marine Geology, to be published.